Toxicology COLLECTION INSTRUCTIONS (TX-FR-0026) VERSION 1.3





MATERIALS PROVIDED

1 Sterile Urine Cup | 1 Urine Transport Tube | 1 Specimen Bag | Lab Pak Mailer

PATIENT PROCEDURE

- 1. Wash hands thoroughly with warm water and soap.
- 2. Open the sterile cup without touching the rim or inner surface of the cup or lid.
 - a. Females: Holding the labia apart, collect first 15-25 ML urine into the provided sterile cup.
 - b. Males: Retract foreskin (if applicable), collect first 15-25 ML urine into the provided sterile cup.
- 3. Securely place cap on urine cup and give to clinical technician.

CLINICAL TECHNICIAN PROCEDURE

- 1. Label the transport tube with two unique patient identifiers. Confirm identity with the patient.
- 2. Put on gloves.
- 3. Do not open urine cup.
- 4. Expose port access on urine cup and aspirate liquid into urine transport tube.
- 5. Place the urine transport tube into the provided specimen bag.
- 6. Remove gloves and perform hand hygiene.
- 7. Place all samples collected for the day into a provided Lab Pak mailer. Seal the mailer and place a shipping label on the outside.

7650 Hub Parkway
Valley View, OH 44125

合 CLIA: #36D2281647 CAP: #9665545